arXiv:0903.4604v1 [math.RA] 26 Mar 2009

THE CLASSIFICATION OF LEIBNIZ SUPERALGEBRAS OF NILINDEX $n+m$ $(m \neq 0.)$

J. R. GÓMEZ, A.KH. KHUDOYBERDIYEV AND B.A. OMIROV

Abstract. In this paper we investigate the description of the complex Leibniz superalgebras with nilindex $n+m$, where n and m $(m \neq 0)$ are dimensions of even and odd parts, respectively. In fact, such superalgebras with characteristic sequence equal to $(n_1, \dots, n_k | m_1, \dots, m_s)$ (where $n_1 + \dots + n_k = n$, $m_1 + \dots + m_s = m$) for $n_1 \geq n-1$ and $(n_1, \dots, n_k | m)$ were classified in works [3]–[6]. Here we prove that in the case of $(n_1, \dots, n_k | m_1, \dots, m_s)$, where $n_1 \leq n-2$ and $m_1 \leq m-1$ the Leibniz superalgebras have nilindex less than $n+m$. Thus, we complete the classification of Leibniz superalgebras with nilindex $n+m$.

Mathematics Subject Classification 2000: 17A32, 17B30, 17B70, 17A70.

Key Words and Phrases: Lie superalgebras, Leibniz superalgebras, nilindex, characteristic sequence, natural gradation.

1. Introduction

During many years the theory of Lie superalgebras has been actively studied by many mathematicians and physicists. A systematic exposition of basic of Lie superalgebras theory can be found in [9]. Many works have been devoted to the study of this topic, but unfortunately most of them do not deal with nilpotent Lie superalgebras. In works [2], [7], [8] the problem of the description of some classes of nilpotent Lie superalgebras have been studied. It is well known that Lie superalgebras are a generalization of Lie algebras. In the same way, the notion of Leibniz algebras, which were introduced in [11], can be generalized to Leibniz superalgebras [1], [10]. Some elementary properties of Leibniz superalgebras were obtained in [1].

In the work [8] the Lie superalgebras with maximal nilindex were classified. Such superalgebras are two-generated and its nilindex equal to $n+m$ (where n and m are dimensions of even and odd parts, respectively). In fact, there exists unique Lie superalgebra of maximal nilindex. This superalgebra is filiform Lie superalgebra (the characteristic sequence equal to $(n-1, 1|m)$) and we mention about paper [2], where some crucial properties of filiform Lie superalgebras are given.

For nilpotent Leibniz superalgebras the description of the case of maximal nilindex (nilpotent Leibniz superalgebras distinguished by the feature of being single-generated) is not difficult and was done in [1].

However, the description of Leibniz superalgebras of nilindex $n+m$ is a very problematic one and it needs to solve many technical tasks. Therefore, they can be studied by applying restrictions on their characteristic sequences. In the present paper

---

The first author was supported by the PAI, FQM143 of the Junta de Andalucía (Spain) and the last author was supported by grant NATO-Reintegration ref. CBP.EAP.RIG.983169.

we consider Leibniz superalgebras with characteristic sequence $(n_1, \dots, n_k|m_1, \dots, m_s)$ $(n_1 \leq n-2$ and $m_1 \leq m-1)$ and nilindex $n+m$. Recall, that such superalgebras for $n_1 \geq n-1$ or $m_1 = m$ have been already classified in works [3]–[6]. Namely, we prove that a Leibniz superalgebra with characteristic sequence equal to $(n_1, \dots, n_k|m_1, \dots, m_s)$ $(n_1 \leq n-2$ and $m_1 \leq m-1)$ has nilindex less than $n+m$. Therefore, we complete classification of Leibniz superalgebras with nilindex $n+m$.

It should be noted that in our study the natural gradation of even part of Leibniz superalgebra played one of the crucial roles. In fact, we used some properties of naturally graded Lie and Leibniz algebras for obtaining the convenience basis of even part of the superalgebra (so-called adapted basis).

Throughout this work we shall consider spaces and (super)algebras over the field of complex numbers. By asterisks $(*)$ we denote the appropriate coefficients at the basic elements of superalgebra.

2. Preliminaries

Recall the notion of Leibniz superalgebras.

Definition 2.1. *A $\mathbb{Z}_2$-graded vector space $L = L_0 \oplus L_1$ is called a Leibniz superalgebra if it is equipped with a product $[-,-]$ which satisfies the following conditions:*

1. $[L_\alpha, L_\beta] \subseteq L_{\alpha+\beta (mod\ 2)}$,

2. $[x,[y,z]] = [[x,y],z] - (-1)^{\alpha\beta}[[x,z],y]-$ Leibniz superidentity,

for all $x \in L$, $y \in L_\alpha$, $z \in L_\beta$ and $\alpha, \beta \in \mathbb{Z}_2$.

The vector spaces L_0 and L_1 are said to be even and odd parts of the superalgebra L, respectively. Evidently, even part of the Leibniz superalgebra is a Leibniz algebra.

Note that if in Leibniz superalgebra L the identity

$$[x,y] = -(-1)^{\alpha\beta}[y,x]$$

holds for any $x \in L_\alpha$ and $y \in L_\beta$, then the Leibniz superidentity can be transformed into the Jacobi superidentity. Thus, Leibniz superalgebras are a generalization of Lie superalgebras and Leibniz algebras.

The set of all Leibniz superalgebras with the dimensions of the even and odd parts, respectively equal to n and m, we denote by $Leib_{n,m}$.

For a given Leibniz superalgebra L we define the descending central sequence as follows:

$$L^1 = L, \quad L^{k+1} = [L^k, L], \quad k \geq 1.$$

Definition 2.2. *A Leibniz superalgebra L is called nilpotent, if there exists $s \in \mathbb{N}$ such that $L^s = 0$. The minimal number s with this property is called nilindex of the superalgebra L.*

Definition 2.3. *The set*

$$\mathcal{R}(L) = \{z \in L \mid [L,z] = 0\}$$

is called the right annihilator of a superalgebra L.

Using the Leibniz superidentity it is easy to see that $\mathcal{R}(L)$ is an ideal of the superalgebra L. Moreover, the elements of the form $[a,b] + (-1)^{\alpha\beta}[b,a]$, $(a \in L_\alpha,\ b \in L_\beta)$ belong to $\mathcal{R}(L)$.

The following theorem describes nilpotent Leibniz superalgebras with maximal nilindex.

Theorem 2.1. [1] *Let L be a Leibniz superalgebra of $Leib_{n,m}$ with nilindex equal to $n+m+1$. Then L is isomorphic to one of the following non-isomorphic superalgebras:*

$$[e_i, e_1] = e_{i+1}, \ 1 \leq i \leq n-1, \ m = 0; \quad \left\{ \begin{array}{ll} [e_i, e_1] = e_{i+1}, & 1 \leq i \leq n+m-1, \\ [e_i, e_2] = 2e_{i+2}, & 1 \leq i \leq n+m-2, \end{array} \right.$$

(omitted products are equal to zero).

Remark 2.1. From the assertion of Theorem 2.1 we have that in case of non-trivial odd part L_1 of the superalgebra L there are two possibility for n and m, namely, $m = n$ if $n+m$ is even and $m = n+1$ if $n+m$ is odd. Moreover, it is clear that the Leibniz superalgebra has the maximal nilindex if and only if it is single-generated.

Let $L = L_0 \oplus L_1$ be a nilpotent Leibniz superalgebra. For an arbitrary element $x \in L_0$, the operator of right multiplication $R_x : L \to L$ (defined as $R_x(y) = [y, x]$) is a nilpotent endomorphism of the space L_i, where $i \in \{0, 1\}$. Taking into account the property of complex endomorphisms we can consider the Jordan form for R_x. For operator R_x denote by $C_i(x)$ $(i \in \{0, 1\})$ the descending sequence of its Jordan blocks dimensions. Consider the lexicographical order on the set $C_i(L_0)$.

Definition 2.4. *A sequence*

$$C(L) = \left(\max_{x \in L_0 \setminus L_0^2} C_0(x) \,\middle|\, \max_{\widetilde{x} \in L_0 \setminus L_0^2} C_1(\widetilde{x}) \right)$$

is said to be the characteristic sequence of the Leibniz superalgebra L.

Similarly to [7] (corollary 3.0.1) it can be proved that the characteristic sequence is invariant under isomorphism.

Since Leibniz superalgebras from $Leib_{n,m}$ with nilindex $n+m$ and with characteristic sequences equal to $(n_1, \dots, n_k | m_1, \dots, m_s)$ either $n_1 \geq n-1$ or $m_1 = m$ were already classified, we shall reduce our investigation to the case of the characteristic sequence $(n_1, \dots, n_k | m_1, \dots m_s)$, where $n_1 \leq n-2$ and $m_1 \leq m-1$

From the Definition 2.4 we have that a Leibniz algebra L_0 has characteristic sequence $(n_1, \dots, n_k)$. Let $l \in \mathbb{N}$ be a nilindex of the Leibniz algebra L_0. Since $n_1 \leq n-2$, then we have $l \leq n-1$ and Leibniz algebra L_0 has at least two generators (the elements which belong to the set $L_0 \setminus L_0^2$).

For the completeness of the statement below we present the classifications of the papers [3]–[5] and [8].

$Leib_{1,m}$:

$$\{ \ [y_i, x_1] = y_{i+1}, \quad 1 \leq i \leq m-1.$$

$Leib_{n,1}$:

$$\left\{ \begin{array}{ll} [x_i, x_1] = x_{i+1}, & 1 \leq i \leq n-1, \\ [y_1, y_1] = \alpha x_n, & \alpha = \{0, \ 1\}. \end{array} \right.$$

$Leib_{2,2}$:

$$\left\{ \begin{array}{l} [y_1, x_1] = y_2, \\ [x_1, y_1] = \frac{1}{2} y_2, \\ [x_2, y_1] = y_2, \\ [y_1, x_2] = 2y_2, \\ [y_1, y_1] = x_2, \end{array} \right. \qquad \left\{ \begin{array}{l} [y_1, x_1] = y_2, \\ [x_2, y_1] = y_2, \\ [y_1, x_2] = 2y_2, \\ [y_1, y_1] = x_2. \end{array} \right.$$

$Leib_{2,m},\ m$ is odd :

$$\left\{\begin{array}{ll} [x_1,x_1]=x_2, & m\geq 3,\\ [y_i,x_1]=y_{i+1}, & 1\leq i\leq m-1,\\ [x_1,y_i]=-y_{i+1}, & 1\leq i\leq m-1,\\ [y_i,y_{m+1-i}]=(-1)^{j+1}x_2, & 1\leq i\leq \frac{m+1}{2}. \end{array}\right. \qquad \left\{\begin{array}{ll} [y_i,x_1]=[x_1,y_i]=-y_{i+1}, & 1\leq i\leq m-1,\\ [y_{m+1-i},y_i]=(-1)^{j+1}x_2, & 1\leq i\leq \frac{m+1}{2}. \end{array}\right.$$

In order to present the classification of Leibniz superalgebras with characteristic sequence $(n-1,1|m)$, $n\geq 3$ and nilindex $n+m$ we need to introduce the following families of superalgebras:

$\mathbf{Leib_{n,n-1}}$:

$L(\alpha_4,\alpha_5,\dots,\alpha_n,\theta)$:

$$\left\{\begin{array}{ll} [x_1,x_1]=x_3, & \\ [x_i,x_1]=x_{i+1}, & 2\leq i\leq n-1,\\ [y_j,x_1]=y_{j+1}, & 1\leq j\leq n-2,\\ [x_1,y_1]=\frac{1}{2}y_2, & \\ [x_i,y_1]=\frac{1}{2}y_i, & 2\leq i\leq n-1,\\ [y_1,y_1]=x_1, & \\ [y_j,y_1]=x_{j+1}, & 2\leq j\leq n-1,\\ [x_1,x_2]=\alpha_4x_4+\alpha_5x_5+\ldots+\alpha_{n-1}x_{n-1}+\theta x_n, & \\ [x_j,x_2]=\alpha_4x_{j+2}+\alpha_5x_{j+3}+\ldots+\alpha_{n+2-j}x_n, & 2\leq j\leq n-2,\\ [y_1,x_2]=\alpha_4y_3+\alpha_5y_4+\ldots+\alpha_{n-1}y_{n-2}+\theta y_{n-1}, & \\ [y_j,x_2]=\alpha_4y_{j+2}+\alpha_5y_{j+3}+\ldots+\alpha_{n+1-j}y_{n-1}, & 2\leq j\leq n-3. \end{array}\right.$$

$G(\beta_4,\beta_5,\dots,\beta_n,\gamma)$:

$$\left\{\begin{array}{ll} [x_1,x_1]=x_3, & \\ [x_i,x_1]=x_{i+1}, & 3\leq i\leq n-1,\\ [y_j,x_1]=y_{j+1}, & 1\leq j\leq n-2,\\ [x_1,x_2]=\beta_4x_4+\beta_5x_5+\ldots+\beta_nx_n, & \\ [x_2,x_2]=\gamma x_n, & \\ [x_j,x_2]=\beta_4x_{j+2}+\beta_5x_{j+3}+\ldots+\beta_{n+2-j}x_n, & 3\leq j\leq n-2,\\ [y_1,y_1]=x_1, & \\ [y_j,y_1]=x_{j+1}, & 2\leq j\leq n-1,\\ [x_1,y_1]=\frac{1}{2}y_2, & \\ [x_i,y_1]=\frac{1}{2}y_i, & 3\leq i\leq n-1,\\ [y_j,x_2]=\beta_4y_{j+2}+\beta_5y_{j+3}+\ldots+\beta_{n+1-j}y_{n-1}, & 1\leq j\leq n-3. \end{array}\right.$$

$\mathbf{Leib_{n,n}}$:

$M(\alpha_4,\alpha_5,\dots,\alpha_n,\theta,\tau)$:

$$\begin{cases}
[x_1,x_1]=x_3, & \\
[x_i,x_1]=x_{i+1}, & 2\le i\le n-1,\\
[y_j,x_1]=y_{j+1}, & 1\le j\le n-1,\\
[x_1,y_1]=\frac{1}{2}y_2, & \\
[x_i,y_1]=\frac{1}{2}y_i, & 2\le i\le n,\\
[y_1,y_1]=x_1, & \\
[y_j,y_1]=x_{j+1}, & 2\le j\le n-1,\\
[x_1,x_2]=\alpha_4x_4+\alpha_5x_5+\ldots+\alpha_{n-1}x_{n-1}+\theta x_n, & \\
[x_2,x_2]=\gamma_4x_4, & \\
[x_j,x_2]=\alpha_4x_{j+2}+\alpha_5x_{j+3}+\ldots+\alpha_{n+2-j}x_n, & 3\le j\le n-2,\\
[y_1,x_2]=\alpha_4y_3+\alpha_5y_4+\ldots+\alpha_{n-1}y_{n-2}+\theta y_{n-1}+\tau y_n, & \\
[y_2,x_2]=\alpha_4y_4+\alpha_5y_4+\ldots+\alpha_{n-1}y_{n-1}+\theta y_n, & \\
[y_j,x_2]=\alpha_4y_{j+2}+\alpha_5y_{j+3}+\ldots+\alpha_{n+2-j}y_n, & 3\le j\le n-2.
\end{cases}$$

$H(\beta_4,\beta_5,\ldots,\beta_n,\delta,\gamma):$

$$\begin{cases}
[x_1,x_1]=x_3, & \\
[x_i,x_1]=x_{i+1}, & 3\le i\le n-1,\\
[y_j,x_1]=y_{j+1}, & 1\le j\le n-2,\\
[x_1,x_2]=\beta_4x_4+\beta_5x_5+\ldots+\beta_nx_n, & \\
[x_2,x_2]=\gamma x_n, & \\
[x_j,x_2]=\beta_4x_{j+2}+\beta_5x_{j+3}+\ldots+\beta_{n+2-j}x_n, & 3\le j\le n-2,\\
[y_1,y_1]=x_1, & \\
[y_j,y_1]=x_{j+1}, & 2\le j\le n-1,\\
[x_1,y_1]=\frac{1}{2}y_2, & \\
[x_i,y_1]=\frac{1}{2}y_i, & 3\le i\le n-1,\\
[y_1,x_2]=\beta_4y_3+\beta_5y_4+\ldots+\beta_ny_{n-1}+\delta y_n, & \\
[y_j,x_2]=\beta_4y_{j+2}+\beta_5y_{j+3}+\ldots+\beta_{n+2-j}y_n, & 2\le j\le n-2.
\end{cases}$$

Analogously, for the Leibniz superalgebras with characteristic sequence $(n|m-1,1)$, $n\geq 2$ we introduce the following families of superalgebras:

$$\mathbf{Leib_{n,n+1}}:$$

$E\left(\gamma,\beta_{\left[\frac{n+4}{2}\right]},\beta_{\left[\frac{n+4}{2}\right]+1},\ldots,\beta_n,\beta\right):$

$$\begin{cases} [x_i, x_1] = x_{i+1}, & 1 \le i \le n-1, \\ [y_j, x_1] = y_{j+1}, & 1 \le j \le n-1, \\ [x_i, y_1] = \frac{1}{2} y_{i+1}, & 1 \le i \le n-1, \\ [y_j, y_1] = x_j, & 1 \le j \le n, \\ [y_{n+1}, y_{n+1}] = \gamma x_n, & \\ [x_i, y_{n+1}] = \sum\limits_{k=\left[\frac{n+4}{2}\right]}^{n+1-i} \beta_k y_{k-1+i}, & 1 \le i \le \left[\frac{n-1}{2}\right], \\ [y_1, y_{n+1}] = -2 \sum\limits_{k=\left[\frac{n+4}{2}\right]}^{n} \beta_k x_{k-1} + \beta x_n, & \\ [y_j, y_{n+1}] = -2 \sum\limits_{k=\left[\frac{n+4}{2}\right]}^{n+2-j} \beta_k x_{k-2+j}, & 2 \le j \le \left[\frac{n+1}{2}\right]. \end{cases}$$

$\mathbf{Leib_{n,n+2}}$:

$F\left(\beta_{\left[\frac{n+5}{2}\right]}, \beta_{\left[\frac{n+5}{2}\right]+1}, \dots, \ \beta_{n+1}\right)$:

$$\begin{cases} [x_i, x_1] = x_{i+1}, & 1 \le i \le n-1, \\ [y_j, x_1] = y_{j+1}, & 1 \le j \le n, \\ [x_i, y_1] = \frac{1}{2} y_{i+1}, & 1 \le i \le n, \\ [y_j, y_1] = x_j, & 1 \le j \le n, \\ [x_i, y_{n+2}] = \sum\limits_{k=\left[\frac{n+5}{2}\right]}^{n+2-i} \beta_k y_{k-1+i}, & 1 \le i \le \left[\frac{n}{2}\right], \\ [y_j, y_{n+2}] = -2 \sum\limits_{k=\left[\frac{n+5}{2}\right]}^{n+2-j} \beta_k x_{k-2+j}, & 1 \le j \le \left[\frac{n}{2}\right] \end{cases}$$

Let us introduce also the following operators which act on k-dimensional vectors:

$$V^0_{j,k}(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, \dots, 0, \overset{j}{1}, \quad \delta \sqrt[j]{\delta^{j+1}} S^{j+1}_{m,j} \alpha_{j+1}, \delta \sqrt[j]{\delta^{j+2}} S^{j+2}_{m,j} \alpha_{j+2}, \dots, \delta \sqrt[j]{\delta^k} S^k_{m,j} \alpha_k);$$

$$V^1_{j,k}(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, \dots, 0, \overset{j}{1}, \quad S^{j+1}_{m,j} \alpha_{j+1}, S^{j+2}_{m,j} \alpha_{j+2}, \dots, S^k_{m,j} \alpha_k);$$

$$V^2_{j,k}(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, \dots, 0, \overset{j}{1}, \quad S^{2(j+1)+1}_{m,2j+1} \alpha_{j+1}, S^{2(j+2)+1}_{m,2j+1} \alpha_{j+2}, \dots, S^{2k+1}_{m,2j+1} \alpha_k);$$

$$V^0_{k+1,k}(\alpha_1, \alpha_2, \dots, \alpha_k) = V^1_{k+1,k}(\alpha_1, \alpha_2, \dots, \alpha_k) = V^2_{k+1,k}(\alpha_1, \alpha_2, \dots, \alpha_k) = (0, 0, \dots, 0);$$

$$W_{s,k}(0, 0, \dots, \overset{j-1}{0}, \overset{j}{1}, S^{j+1}_{m,j} \alpha_{j+1}, S^{j+2}_{m,j} \alpha_{j+2}, \dots, S^k_{m,j} \alpha_k, \gamma) =$$

$$= (0, 0, \dots, \overset{j}{1}, 0, \dots, \overset{s+j}{1}, S^{s+1}_{m,s} \alpha_{s+j+1}, S^{s+2}_{m,s} \alpha_{s+j+2}, \dots, S^{k-j}_{m,s} \alpha_k, S^{k+6-2j}_{m,s} \gamma),$$

$$W_{k+1-j,k}(0, 0, \dots, \overset{j-1}{0}, \overset{j}{1}, S^{j+1}_{m,j} \alpha_{j+1}, S^{j+2}_{m,j} \alpha_{j+2}, \dots, S^k_{m,j} \alpha_k, \gamma) =$$

$$= (0, 0, \dots, \overset{j}{1}, 0, \dots, 1),$$

$$W_{k+2-j,k}(0, 0, \dots, \overset{j-1}{0}, \overset{j}{1}, S^{j+1}_{m,j} \alpha_{j+1}, S^{j+2}_{m,j} \alpha_{j+2}, \dots, S^k_{m,j} \alpha_k, \gamma) =$$

$$= (0, 0, \ldots, \overset{j}{1}, 0, \ldots, 0),$$

where $k \in N$, $\delta = \pm 1$, $1 \leq j \leq k$, $1 \leq s \leq k-j$, and $S_{m,t} = \cos\frac{2\pi m}{t} + i \sin\frac{2\pi m}{t}$ $(m = 0, 1, \ldots, t-1)$.

Below we present the complete list of pairwise non-isomorphic Leibniz superalgebras with $n+m$:

with characteristic sequence equal to $(n-1, 1|m)$:

$$\begin{aligned}
&L\left(V^1_{j,n-3}\left(\alpha_4, \alpha_5, \ldots, \alpha_n\right), S^{n-3}_{m,j}\theta\right), \qquad 1 \leq j \leq n-3,\\
&L(0,0,\ldots,0,1),\ L(0,0,\ldots,0),\ G(0,0,\ldots,0,1),\ G(0,0,\ldots,0),\\
&G\left(W_{s,n-2}\left(V^1_{j,n-3}\left(\beta_4, \beta_5, \ldots, \beta_n\right), \gamma\right)\right), \quad 1 \leq j \leq n-3,\ 1 \leq s \leq n-j,\\
&M\left(V^1_{j,n-2}\left(\alpha_4, \alpha_5, \ldots, \alpha_n\right), S^{n-3}_{m,j}\theta\right), \qquad 1 \leq j \leq n-2,\\
&M(0,0,\ldots,0,1),\ M(0,0,\ldots,0),\ H(0,0,\ldots,0,1),\ H(0,0,\ldots,0),\\
&H\left(W_{s,n-1}\left(V^1_{j,n-2}\left(\beta_4, \beta_5, \ldots, \beta_n\right), \gamma\right)\right), \quad 1 \leq j \leq n-2,\ 1 \leq s \leq n+1-j,
\end{aligned}$$

with characteristic sequence equal to $(n|m-1, 1)$ if n is odd (i.e. $n = 2q-1$):

$$\begin{aligned}
&E\left(1, \delta\beta_{q+1}, V^0_{j,q-2}(\beta_{q+2}, \beta_{q+3}, \ldots, \beta_n), 0\right), && \beta_{q+1} \neq \pm\frac{1}{2}, && 1 \leq j \leq q-1,\\
&E\left(1, \beta_{q+1}, V^0_{j,q-1}(\beta_{q+2}, \beta_{q+3}, \ldots, \beta_n, \beta)\right), && \beta_{q+1} = \pm\frac{1}{2}, && 1 \leq j \leq q,\\
&E(0, 1, V^0_{j,q-2}(\beta_{q+2}, \beta_{q+3}, \ldots, \beta_n), 0), && 1 \leq j \leq q-1, &&\\
&E(0, 0, W_{s,q-1}(V^1_{j,q-1}(\beta_{q+2}, \beta_{q+3}, \ldots, \beta_n, \beta))), && 1 \leq j \leq q-1, && 1 \leq s \leq q-j,\\
&E(0, 0, \ldots, 0); &&&&
\end{aligned}$$

if n is even (i.e. $n = 2q$):

$$\begin{aligned}
&E(1, V^2_{j,q-1}(\beta_{q+2}, \beta_{q+3}, \ldots, \beta_n,), 0), && 1 \leq j \leq q,\\
&E(0, W_{s,q}(V^1_{j,q}(\beta_{q+2}, \beta_{q+3}, \ldots, \beta_n, \beta))), && 1 \leq j \leq q, \quad 1 \leq s \leq q+1-j,\\
&E(0, 0, \ldots, 0). &&
\end{aligned}$$

$$F\left(W_{s,n+2-\left[\frac{n+5}{2}\right]}\left(V^1_{j,n+2-\left[\frac{n+5}{2}\right]}\left(\beta_{\left[\frac{n+5}{2}\right]}, \beta_{\left[\frac{n+5}{2}\right]+1}, \ldots, \beta_{n+1}\right)\right)\right),$$

where $1 \leq j \leq n+2-\left[\dfrac{n+5}{2}\right]$, $1 \leq s \leq n+3-\left[\dfrac{n+5}{2}\right]-j$,

$$F(0, 0, \ldots, 0).$$

For a given Leibniz algebra A of the nilindex l we put $gr(A)_i = A^i/A^{i+1}$, $1 \leq i \leq l-1$ and $gr(A) = gr(A)_1 \oplus gr(A)_2 \oplus \cdots \oplus gr(A)_{l-1}$. Then $[gr(A)_i, gr(A)_j] \subseteq gr(A)_{i+j}$ and we obtain the graded algebra $gr(A)$.

Definition 2.5. *The gradation constructed in this way is called the natural gradation and if a Leibniz algebra G is isomorphic to $gr(A)$ we say that the algebra G is naturally graded Leibniz algebra.*

3. The main result

Let L be a Leibniz superalgebra with characteristic sequence $(n_1, \dots, n_k|m_1, \dots, m_s)$, where $n_1 \leq n-2$, $m_1 \leq m-1$ and of nilindex $n+m$. Since the second part of the characteristic sequence of the Leibniz superalgebra L is equal to $(m_1, \dots, m_s)$ then by the definition of the characteristic sequence there exists a nilpotent endomorphism R_x ($x \in L_0 \setminus L_0^2$) of the space L_1 such that its Jordan form consists of s Jordan blocks. Therefore, we can assume the existence of an adapted basis $\{y_1, y_2, \dots, y_m\}$ of the subspace L_1, such that

$$\left\{ \begin{array}{ll} [y_j, x] = y_{j+1}, & j \notin \{m_1, m_1+m_2, \dots, m_1+m_2+\dots+m_s\}, \\ [y_j, x] = 0, & j \in \{m_1, m_1+m_2, \dots, m_1+m_2+\dots+m_s\}. \end{array} \right. \tag{1}$$

for some $x \in L_0 \setminus L_0^2$.

Further we shall use a homogeneous basis $\{x_1, \dots, x_n\}$ with respect to natural gradation of the Leibniz algebra L_0, which is also agreed with the lower central sequence of L.

The main result of the paper is that the nilindex of the Leibniz superalgebra L with characteristic sequence $(n_1, \dots, n_k|m_1, \dots, m_s)$, $n_1 \leq n-2, m_1 \leq m-1$ is less than $n+m$.

According to the Theorem 2.1 we have the description of single-generated Leibniz superalgebras, which have nilindex $n+m+1$. If the number of generators is greater than two, then superalgebra has nilindex less than $n+m$. Therefore, we should consider case of two-generated superalgebras.

The possible cases for the generators are:

1. Both generators lie in L_0, i.e. $dim(L^2)_0 = n-2$ and $dim(L^2)_1 = m$;
2. One generator lies in L_0 and another one lies in L_1, i.e. $dim(L^2)_0 = n-1$ and $dim(L^2)_1 = m-1$;
3. Both generators lie in L_1, i.e. $dim(L^2)_0 = n$ and $dim(L^2)_1 = m-2$.

Moreover, two-generated superalgebra L has nilindex $n+m$ if and only if $dim L^k = n+m-k$ for $2 \leq k \leq n+m$.

Since $m \neq 0$ we omit the case where both generators lie in even part.

3.1. **The case of one generator in L_0 and another one in L_1.**

Since $dim(L^2)_0 = n-1$ and $dim(L^2)_1 = m-1$ then there exist some m_j, $0 \leq j \leq s-1$ (here we assume $m_0 = 0$) such that $y_{m_1+\dots+m_j+1} \notin L^2$. By a shifting of basic elements we can assume that $m_j = m_0$, i.e. the basic element y_1 can be chosen as a generator of the superalgebra L. Of course, by this shifting the condition from definition of the characteristic sequence $m_1 \geq m_2 \geq \dots \geq m_s$ can be broken, but further we shall not use the condition.

Let $L = L_0 \oplus L_1$ be a two generated Leibniz superalgebra from $Leib_{n,m}$ with characteristic sequence equal to $(n_1, \dots, n_k|m_1, \dots, m_s)$ and let $\{x_1, \dots, x_n, y_1, \dots, y_m\}$ be a basis of the L.

Lemma 3.1. *Let one generator lies in L_0 and another one lies in L_1. Then x_1 and y_1 can be chosen as generators of the L. Moreover, in equality (1) instead of element x we can suppose x_1.*

Proof. As mentioned above y_1 can be chosen as the first generator of L. If $x \in L \setminus L^2$ then the assertion of the lemma is evident. If $x \in L^2$ then there exists some i_0 ($2 \leq i_0$) such that $x_{i_0} \in L \setminus L^2$. Set $x_1' = Ax + x_{i_0}$ for $A \neq 0$ then x_1' is a generator of the superalgebra L (since $x_1' \in L \setminus L^2$). Moreover, making transformation of the basis of L_1 as follows

$$\begin{cases} y_j' = y_j, & j \in \{1, m_1+1, \dots, m_1+m_2+\dots+m_{s-1}+1\}, \\ y_j' = [y_{j-1}', x_1'], & j \notin \{1, m_1+1, \dots, m_1+m_2+\dots+m_{s-1}+1\}. \end{cases}$$

and taking sufficiently big value of the parameter A we preserve the equality (1). Thus, in the basis $\{x_1', x_2, \dots, x_n, y_1', y_2', \dots, y_m'\}$ the elements x_1' and y_1' are generators. □

Due to Lemma 3.1 further we shall suppose that $\{x_1, y_1\}$ are generators of the Leibniz superalgebra L. Therefore,

$$L^2 = \{x_2, x_3, \dots, x_n, y_2, y_3, \dots, y_m\}.$$

Let us introduce the notations:

$$[x_i, y_1] = \sum_{j=2}^{m} \alpha_{i,j} y_j, \ 1 \leq i \leq n, \quad [y_i, y_1] = \sum_{j=2}^{n} \beta_{i,j} x_j, \ 1 \leq i \leq m. \tag{2}$$

Without loss of generality we can assume that $y_{m_1+\dots+m_i+1} \in L^{t_i} \setminus L^{t_i+1}$, where $t_i < t_j$ for $1 \leq i < j \leq s-1$.

Firstly we consider the case of $dim(L^3)_0 = n-1$, then $dim(L^3)_0 = n-2$.

Case $dim(L^3)_0 = n-1$.

In this subcase we have

$$L^3 = \{x_2, x_3, \dots, x_n, y_3, \dots, y_{m_1}, B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+2}, \dots, y_m\},$$

where $(B_1, B_2) \neq (0,0)$.

Analyzing the way the element x_2 can be obtained, we conclude that there exist i_0 $(2 \leq i_0 \leq m)$ such that $[y_{i_0}, y_1] = \sum\limits_{j=2}^{n} \beta_{i_0,2} x_j, \ \beta_{i_0,2} \neq 0$.

Let us show that $i_0 \notin \{m_1+1, \dots, m_1+\dots+m_{s-1}+1\}$. It is known that the elements $y_{m_1+m_2+1}, \dots, y_{m_1+\dots+m_{s-1}+1}$ are generated from the products $[x_i, y_1]$, $(2 \leq i \leq n)$. Due to nilpotency of L we get $i_0 \notin \{m_1+m_2+1, \dots, m_1+\dots+m_{s-1}+1\}$. If y_{m_1+1} is generated by $[x_1, y_1]$, i.e. in the expression $[x_1, y_1] = \sum\limits_{j=2}^{m} \alpha_{1,j} y_j \ \alpha_{1,m_1+1} \neq 0$ then we consider the product

$$[[x_1, y_1], y_1] = [\sum_{j=2}^{m} \alpha_{1,j} y_j, y_1] = \alpha_{1,m_1+1} \beta_{m_1+1,2} x_2 + \sum_{i \geq 3} (*) x_i.$$

On the other hand,

$$[[x_1, y_1], y_1] = \frac{1}{2}[x_1, [y_1, y_1]] = \frac{1}{2}[x_1, \sum_{j=2}^{n} \beta_{1,j} x_j] = \sum_{i \geq 3} (*) x_i$$

Comparing the coefficients at the corresponding basic elements we obtain $\alpha_{1,m_1+1}\beta_{m_1+1,2} = 0$, which implies $\beta_{m_1+1,2} = 0$. It means that $i_0 \neq m_1+1$. Therefore, $\beta_{i_0,2} \neq 0$, where $i_0 \notin \{m_1+1, \dots, m_1+\dots+m_{s-1}+1\}$.

Case $y_2 \notin L^3$. Then $B_2 \neq 0$. Let $h \in \mathbb{N}$ be a number such that $x_2 \in L^h \setminus L^{h+1}$, that is

$$L^h = \{x_2, x_3, \dots, x_n, y_h, \dots, y_{m_1}, B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+2}, \dots, y_m\}, \ h \geq 3,$$

$$L^{h+1} = \{x_3, x_4, \dots, x_n, y_h, \dots, y_{m_1}, B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+2}, \dots, y_m\}.$$

Since the elements $B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+m_2+1}, \dots, y_{m_1+\dots+m_{s-1}+1}$ are generated from the multiplications $[x_i, y_1], 2 \leq i \leq n$ it follows that $h \leq m_1 + 1$.

So, x_2 can be obtained only from product $[y_{h-1}, y_1]$ and thereby $\beta_{h-1,2} \neq 0$. Making the change $x_2' = \sum\limits_{j=2}^{n} \beta_{h-1,j} x_j$ we can assume that $[y_{h-1}, y_1] = x_2$.

Let now p is a number such that $y_h \in L^{h+p} \setminus L^{h+p+1}$. Then for the powers of superalgebra L we have the following

$$L^{h+p} = \{x_{p+2}, x_{p+3}, \dots, x_n, y_h, \dots, y_{m_1}, B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+2}, \dots, y_m\}, \ p \geq 1,$$

$$L^{h+p+1} = \{x_{p+2}, x_{p+3}, \dots, x_n, y_{h+1}, \dots, y_{m_1}, B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+2}, \dots, y_m\}.$$

In the following lemma the useful expression for the products $[y_i, y_j]$ is presented.

Lemma 3.2. *The equality:*

$$[y_i, y_j] = (-1)^{h-1-i} C_{j-1}^{h-1-i} x_{i+j+2-h} + \sum_{t > i+j+2-h} (*) x_t, \tag{3}$$

$1 \leq i \leq h-1, \ h-i \leq j \leq min\{h-1, h-1+p-i\}$, *holds.*

Proof. The proof is deduced by the induction on j at any value of i. □

For the natural number p we have the following

Lemma 3.3. *Under the above conditions* $p = 1$.

Proof. Assume the contrary, i.e. $p > 1$. Then we can suppose

$$[x_i, x_1] = x_{i+1}, \ 2 \leq i \leq p, \quad [x_{p+1}, y_1] = \sum_{j=h}^{m} \alpha_{p+1,j} y_j, \quad \alpha_{p+1,h} \neq 0.$$

Using the equality (3) we consider the following chain of equalities

$$[y_1, [y_{h-1}, x_1]] = [[y_1, y_{h-1}], x_1] - [[y_1, x_1], y_{h-1}] = (-1)^{h-2} x_3 + \sum_{t \geq 4} (*) x_t -$$

$$-(-1)^{h-3}(h-2) x_3 + \sum_{t \geq 4} (*) x_t = (-1)^h (h-1) x_3 + \sum_{t \geq 4} (*) x_t.$$

If $h \leq m_1$, then $[y_1, [y_{h-1}, x_1]] = [y_1, y_h]$. Since $y_h \in L^{h+p}$ and $p > 1$ then in the decomposition of $[y_1, y_h]$ the coefficient at the basic elements x_2 and x_3 are equal to zero. Therefore, from the above equalities we get a contradiction with assumption $p > 1$.

If $h = m_1 + 1$, then $[y_1, [y_{h-1}, x_1]] = 0$ and we also obtain the irregular equality $(-1)^h (h-1) x_3 + \sum\limits_{t \geq 4} (*) x_t = 0$. Therefore, the proof of the lemma is completed. □

We resume our main result in considered cases in the following

Theorem 3.1. *Let $L = L_0 \oplus L_1$ be a Leibniz superalgebra from $Leib_{n,m}$ with characteristic sequence equal to $(n_1, \dots, n_k | m_1, \dots, m_s)$, where $n_1 \leq n-2$, $m_1 \leq m-1$ and let $dim(L^3)_0 = n-1$ with $y_2 \notin L^3$. Then L has a nilindex less than $n+m$.*

Proof. Let us assume the contrary, i.e. nilindex of the superalgebra L equal to $n+m$. Then according to the Lemma 3.3 we have

$$L^{h+2} = \{x_3, \dots, x_n, y_{h+1}, \dots, y_{m_1}, B_1 y_2 + B_2 y_{m_1+1}, y_{m_1+2}, \dots, y_m\}.$$

Since $y_h \notin L^{h+2}$, it follows that

$$\alpha_{2,h} \neq 0, \quad \alpha_{i,h} = 0 \quad \text{for} \quad i > 2.$$

Consider the product

$$[[y_{h-1}, y_1], y_1] = \frac{1}{2}[y_{h-1}, [y_1, y_1]] = \frac{1}{2}[y_{h-1}, \sum_{i=2}^{n} \beta_{1,i} x_i].$$

The element y_{h-1} belongs to L^{h-1} and elements $x_2, x_3, \dots, x_n$ lie in L^3. Hence $\frac{1}{2}[y_{h-1}, \sum\limits_{i=2}^{n} \beta_{1,i} x_i] \in L^{h+2}$. Since $y_h \notin L^{h+2}$, we obtain that $[[y_{h-1}, y_1], y_1] = \sum\limits_{j \geq h+1} (*) y_j$.

On the other hand,

$$[[y_{h-1}, y_1], y_1] = [x_2, y_1] = \alpha_{2,h} y_h + \sum_{j=h+1}^{m} \alpha_{2,j} y_j.$$

Comparing the coefficients at the basic elements we obtain $\alpha_{2,h} = 0$, which is a contradiction with the assumption that the superalgebra L has nilindex equal to $n+m$ and therefore the assertion of the theorem is proved. □

Case $y_2 \in L^3$. Then $B_2 = 0$ and the following theorem is true.

Theorem 3.2. *Let $L = L_0 \oplus L_1$ be a Leibniz superalgebra from $Leib_{n,m}$ with characteristic sequence equal to $(n_1, \dots, n_k | m_1, \dots, m_s)$, where $n_1 \leq n-2$, $m_1 \leq m-1$ and let $dim(L^3)_0 = n-1$ with $y_2 \in L^3$. Then L has a nilindex less than $n+m$.*

Proof. We shall prove the assertion of the theorem by contrary method, i.e. we assume that nilindex of the superalgebra L equal to $n+m$. The condition $y_2 \in L^3$ implies

$$L^3 = \{x_2, x_3, \dots, x_n, y_2, \dots, y_{m_1}, y_{m_1+2}, \dots, y_m\}.$$

Then $\alpha_{1,m_1+1} \neq 0$ and $\alpha_{i,m_1+1} = 0$ for $i \geq 2$. The element y_2 is generated from products $[x_i, y_1]$, $i \geq 2$ which implies $y_2 \in L^4$. Since $[y_{m_1+1}, y_1] = [[x_1, y_1], y_1] = \frac{1}{2}[x_1, [y_1, y_1]] = \frac{1}{2}[x_1, \sum (*) x_i]$ and x_2 is a generator of the Leibniz algebra L_0 then x_2 can not generated from the product $[y_{m_1+1}, y_1]$. Thereby x_2 also belongs to L^4.

Consider the equality

$$[[x_1, y_1], x_1] = [x_1, [y_1, x_1]] + [[x_1, x_1], y_1] = [x_1, y_2] - [\sum_{i \geq 3} (*) x_i, y_1].$$

From this it follows that the product $[[x_1, y_1], x_1]$ belongs to L^5 (and therefore belongs to L^4).

On the other hand,

$$[[x_1, y_1], x_1] = [\sum_{j=2}^{m} \alpha_{1,j} y_j, x_1] = \alpha_{1,2} y_3 + \cdots + \alpha_{1,m_1-1} y_{m_1} + \alpha_{1,m_1+1} y_{m_1+2} + \cdots + \alpha_{1,m-1} y_m.$$

Since $\alpha_{1,m_1+1} \neq 0$, we obtain that $y_{m_1+2} \in L^4$. Thus, we have $L^4 = \{x_2, x_3, \dots, x_n, y_2, \dots, y_{m_1}, y_{m_1+2}, \dots, y_m\}$, that is $L^4 = L^3$. It is a contradiction to nilpotency of the superalgebra L.

Thus, we get a contradiction with assumption that the superalgebra L has nilindex equal to $n+m$ and therefore the assertion of the theorem is proved. □

From Theorems 3.1 and 3.2 we obtain that Leibniz superalgebra L with condition $dim(L^3)_0 = n-1$ has nilindex less than $n+m$.

The investigation of the Leibniz superalgebra with property $dim(L^3)_0 = n-2$ shows that the restriction to nilindex depends on the structure of the Leibniz algebra L_0. Below we present some necessary remarks on nilpotent Leibniz algebras.

Let $A = \{z_1, z_2, \dots, z_n\}$ be an n-dimensional nilpotent Leibniz algebra of nilindex l $(l < n)$. Note that algebra A is not single-generated.

Proposition 3.1. [6] *Let $gr(A)$ be a naturally graded non-Lie Leibniz algebra. Then $dimA^3 \leq n-4$.*

The result on nilindex of the superalgebra under the condition $dim(L^3)_0 = n-2$ is established in the following two theorems.

Theorem 3.3. *Let $L = L_0 \oplus L_1$ be a Leibniz superalgebra from $Leib_{n,m}$ with characteristic sequence $(n_1, \dots, n_k | m_1, \dots m_s)$, where $n_1 \leq n-2$, $m_1 \leq m-1$, $dim(L^3)_0 = n-2$ and $dimL_0^3 \leq n-4$. Then L has a nilindex less than $n+m$.*

Proof. Let us assume the contrary, i.e. the nilindex of the superalgebra L is equal to $n+m$. According to the condition $dim(L^3)_0 = n-2$ we have

$$L^3 = \{x_3, x_4, \dots, x_n, y_2, y_3, \dots, y_m\}.$$

From the condition $dimL_0^3 \leq n-4$ it follows that there exist at least two basic elements, that do not belong to L_0^3. Without loss of generality, one can assume $x_3, x_4 \notin L_0^3$.

Let h be a natural number such that $x_3 \in L^{h+1} \setminus L^{h+2}$, then we have

$$L^{h+1} = \{x_3, x_4, \dots, x_n, y_h, y_{h+1}, \dots, y_m\}, \ h \geq 2, \ \beta_{h-1,3} \neq 0.$$

$$L^{h+2} = \{x_4, \dots, x_n, y_h, y_{h+1}, \dots, y_m\}.$$

Let us suppose $x_3 \notin L_0^2$. Then we have that x_3 can not be obtained by the products $[x_i, x_1]$, with $2 \leq i \leq n$. Therefore, it is generated by products $[y_j, y_1]$, $2 \leq j \leq m$, which implies $h \geq 3$ and $\alpha_{2,2} \neq 0$.

If $h = 3$, then $\beta_{2,3} \neq 0$.

Consider the chain of equalities

$$[[x_2, y_1], y_1] = [\sum_{j=2}^{m} \alpha_{2,j} y_j, y_1] = \sum_{j=2}^{m} \alpha_{2,j} [y_j, y_1] = \alpha_{2,2}\beta_{2,3} x_3 + \sum_{i\geq 4} (*) x_i.$$

On the other hand,

$$[[x_2, y_1], y_1] = \frac{1}{2}[x_2, [y_1, y_1]] = \frac{1}{2}[x_2, \sum_{i=2}^{n} \beta_{1,i} x_i] = \frac{1}{2}\sum_{i=2}^{n} \beta_{1,i} [x_2, x_i] = \sum_{i\geq 4} (*) x_i.$$

Comparing the coefficients at the corresponding basic elements, we get a contradiction with $\beta_{2,3}=0$. Thus, $h \geq 4$.

Since $y_2 \in L^3$ and $h \geq 4$ we have $y_{h-2} \in L^{h-1}$, which implies $[y_{h-2}, y_2] \in L^{h+2} = \{x_4, \dots, x_n, y_h, y_{h+1}, \dots, y_m\}$. It means that in the decomposition $[y_{h-2}, y_2]$ the coefficient at the basic element x_3 is equal to zero.

On the other hand,

$$[y_{h-2}, y_2] = [y_{h-2}, [y_1, x_1]] = [[y_{h-2}, y_1], x_1] - [[y_{h-2}, x_1], y_1] =$$

$$= [\sum_{i=2}^{n} \beta_{h-2,i} x_i, x_1] - [y_{h-1}, y_1] = -\beta_{h-1,3} x_3 + \sum_{i \geq 4} (*) x_i.$$

Hence, we get $\beta_{h-1,3}=0$, which is obtained from the assumption $x_3 \notin L_0^2$.

Therefore, we have $x_3, x_4 \in L_0^2 \setminus L_0^3$. The condition $x_4 \notin L_0^3$ deduce that x_4 can not be obtained by the products $[x_i, x_1]$, with $3 \leq i \leq n$. Therefore, it is generated by products $[y_j, y_1], h \leq j \leq m$. Hence, $L^{h+3} = \{x_4, \dots, x_n, y_{h+1}, \dots, y_m\}$ and $y_h \in L^{h+2} \setminus L^{h+3}$, which implies $\alpha_{3,h} \neq 0$.

Let p $(p \geq 3)$ be a natural number such that $x_4 \in L^{h+p} \setminus L^{h+p+1}$.

Suppose that $p=3$. Then $\beta_{h,4} \neq 0$.

Consider the chain of equalities

$$[[x_3, y_1], y_1] = [\sum_{j=h}^{m} \alpha_{3,j} y_j, y_1] = \sum_{j=h}^{m} \alpha_{3,j} [y_j, y_1] = \alpha_{3,h} \beta_{h,4} x_4 + \sum_{i \geq 5} (*) x_i.$$

On the other hand,

$$[[x_3, y_1], y_1] = \frac{1}{2}[x_3, [y_1, y_1]] = \frac{1}{2}[x_3, \sum_{i=2}^{n} \beta_{1,i} x_i] = \frac{1}{2} \sum_{i=2}^{n} \beta_{1,i} [x_3, x_i] = \sum_{i \geq 5} (*) x_i.$$

Comparing the coefficients at the corresponding basic elements in these equations we get $\alpha_{3,h}\beta_{h,4}=0$, which implies $\beta_{h,4}=0$. It is a contradiction with assumption $p=3$. Therefore, $p \geq 4$ and for the powers of descending lower sequences we have

$$\begin{aligned}
L^{h+p-2} &= \{x_4, \dots, x_n, y_{h+p-4}, \dots, y_m\}, \\
L^{h+p-1} &= \{x_4, \dots, x_n, y_{h+p-3}, \dots, y_m\}, \\
L^{h+p} &= \{x_4, \dots, x_n, y_{h+p-2}, \dots, y_m\}, \\
L^{h+p+1} &= \{x_5, \dots, x_n, y_{h+p-2}, \dots, y_m\}.
\end{aligned}$$

It is easy to see that in the decomposition $[y_{h+p-3}, y_1] = \sum\limits_{i=4}^{n} \beta_{h+p-3,i} x_i$ we have $\beta_{h+p-3,4} \neq 0$.

Consider the equalities

$$[y_{h+p-4}, y_2] = [y_{h+p-4}, [y_1, x_1]] = [[y_{h+p-4}, y_1], x_1] - [[y_{h+p-4}, x_1], y_1] =$$

$$= [\sum_{i=4}^{n} \beta_{h+p-3,i} x_i, x_1] - [y_{h+p-3}, y_1] = -\beta_{h+p-3,4} x_4 + \sum_{i \geq 5} (*) x_i.$$

Since $y_{h+p-4} \in L^{h+p-2}, y_2 \in L^3$ and $\beta_{h+p-3,4} \neq 0$, then the element x_4 should lie in L^{h+p+1}, but it contradicts to $L^{h+p+1} = \{x_5, \dots, x_n, y_{h+p-2}, \dots, y_m\}$. Thus, the superalgebra L has a nilindex less than $n+m$. $\square$

From Theorem 3.3 we conclude that Leibniz superalgebra $L = L_0 \oplus L_1$ with the characteristic sequence $(n_1, \dots, n_k | m_1, \dots, m_s)$, where $n_1 \leq n-2,\ m_1 \leq m-1$ and nilindex $n+m$ can appear only if $dim L_0^3 \geq n-3$. Taking into account the condition $n_1 \leq n-2$ and properties of naturally graded subspaces $gr(L_0)_1,\ gr(L_0)_2$ we get $dim L_0^3 = n-3$.

Let $dim L_0^3 = n-3$. Then

$$gr(L_0)_1 = \{\overline{x}_1, \overline{x}_2\},\ gr(L_0)_2 = \{\overline{x}_3\}.$$

From Proposition 3.1 the naturally graded Leibniz algebra $gr(L_0)$ is a Lie algebra, i.e. the following multiplication rules hold

$$\begin{cases} [\overline{x}_1, \overline{x}_1] = 0, \\ [\overline{x}_2, \overline{x}_1] = \overline{x}_3, \\ [\overline{x}_1, \overline{x}_2] = -\overline{x}_3, \\ [\overline{x}_2, \overline{x}_2] = 0. \end{cases}$$

Using these products for the corresponding products in the Leibniz algebra L_0 with the basis $\{x_1, x_2, \dots, x_n\}$ we have

$$\begin{cases} [x_1, x_1] = \gamma_{1,4}x_4 + \gamma_{1,5}x_5 + \dots + \gamma_{1,n}x_n, \\ [x_2, x_1] = x_3, \\ [x_1, x_2] = -x_3 + \gamma_{2,4}x_4 + \gamma_{2,5}x_5 + \dots + \gamma_{2,n}x_n, \\ [x_2, x_2] = \gamma_{3,4}x_4 + \gamma_{3,5}x_5 + \dots + \gamma_{3,n}x_n. \end{cases} \tag{4}$$

Theorem 3.4. *Let* $L = L_0 \oplus L_1$ *be a Leibniz superalgebra from* $Leib_{n,m}$ *with characteristic sequence* $(n_1, \dots, n_k | m_1, \dots, m_s)$, *where* $n_1 \leq n-2,\ m_1 \leq m-1$, $dim(L^3)_0 = n-2$ *and* $dim L_0^3 = n-3$. *Then* L *has a nilindex less than* $n+m$.

Proof. Let us suppose the contrary, i.e. the nilindex of the superalgebra L equals $n+m$. Then from the condition $dim(L^3)_0 = n-2$ we obtain

$$L^2 = \{x_2, x_3, \dots, x_n, y_2, \dots, y_m\},$$

$$L^3 = \{x_3, x_4, \dots, x_n, y_2, \dots, y_m\}.$$

$$L^4 \supset \{x_4, \dots, x_n, y_3, \dots, y_{m_1}, B_1y_2 + B_2y_{m_1+1}, y_{m_1+2} \dots, y_m\}, \quad (B_1, B_2) \neq (0,0).$$

Suppose $x_3 \notin L^4$. Then

$$L^4 = \{x_4, \dots, x_n, y_2, \dots, y_{m_1}, y_{m_1+1}, \dots, y_m\}.$$

Let $B_1'y_2 + B_2'y_{m_1+1}$ be an element which earlier disappear in the descending lower sequence for L. Then this element can not to be generated from the products $[x_i, y_1],\ 2 \leq i \leq n$. Indeed, since $x_3 \notin L^4$, the element can not to be generated from $[x_2, y_1]$. Due to structure of L_0 the elements $x_i, (3 \leq i \leq n)$ are in L_0^2, i.e. they are generated by the linear combinations of the products of elements from L_0. The equalities

$$[[x_i, x_j], y_1] = [x_i, [x_j, y_1]] + [[x_i, y_1], x_j] = [x_i, \sum_{t=2}^{m} \alpha_{j,t}y_t] + [\sum_{t=2}^{m} \alpha_{i,t}y_t, x_i]$$

derive that the element $B_1'y_2 + B_2'y_{m_1+1}$ can not be obtained by the products $[x_i, y_1], 3 \leq i \leq n$. However, it means that $x_3 \in L^4$. Thus, we have

$$L^4 = \{x_3, x_4, \dots, x_n, y_3, \dots, y_{m_1}, B_1y_2 + B_2y_{m_1+1}, y_{m_1+2}, \dots, y_m\},$$

where $(B_1, B_2) \neq (0,0)$ and $B_1B_2' - B_2B_1' \neq 0$.

The simple analysis of descending lower sequences L^3 and L^4 implies

$$[x_2, y_1] = \alpha'_{2,2}(B'_1y_2+B'_2y_{m_1+1})+\alpha'_{2,m_1+1}(B_1y_2+B_2y_{m_1+1})+\sum_{\substack{j=3\\ j\neq m_1+1}}^{m}\alpha_{2,j}y_j, \quad \alpha'_{2,2}\neq 0.$$

Let h be a natural number such that $x_3 \in L^{h+1} \setminus L^{h+2}$, i.e.

$$L^h = \{x_3, x_4, \dots, x_n, y_{h-1}, y_h, \dots, y_{m_1}, B_1y_2 + B_2y_{m_1+1}, y_{m_1+2}, \dots, y_m\}, h \geq 3,$$

$$L^{h+1} = \{x_3, x_4, \dots, x_n, y_h, y_{h+1}, \dots, y_{m_1}, B_1y_2 + B_2y_{m_1+1}, y_{m_1+2}, \dots, y_m\},$$

$$L^{h+2} = \{x_4, \dots, x_n, y_h, y_{h+1}, \dots, y_{m_1}, B_1y_2 + B_2y_{m_1+1}, y_{m_1+2}, \dots, y_m\}.$$

If $h = 3$, then $[B'_1y_2+B'_2y_{m_1+1}, y_1] = \beta'_{2,3}x_3+\sum\limits_{i\geq 4}(*)x_4$, $\beta'_{2,3}\neq 0$ and we consider the product

$$[[x_2, y_1], y_1] = [\alpha'_{2,2}(B'_1y_2+B'_2y_{m_1+1})+\alpha'_{2,m_1+1}(B_1y_2+B_2y_{m_1+1})+\sum_{\substack{j=3\\ j\neq m_1+1}}^{m}\alpha_{2,j}y_j, y_1] =$$

$$= \alpha'_{2,2}[B'_1y_2 + B'_2y_{m_1+1}, y_1] + \alpha'_{2,m_1+1}[B_1y_2 + B_2y_{m_1+1}, y_1]+$$

$$+\sum_{\substack{j=3\\ j\neq m_1+1}}^{m}\alpha_{2,j}[y_j, y_1] = \alpha'_{2,2}\beta'_{2,3}x_3 + \sum_{i\geq 4}(*)x_4.$$

On the other hand, due to (4) we have

$$[[x_2, y_1], y_1] = \frac{1}{2}[x_2, [y_1, y_1]] = \frac{1}{2}[x_2, \sum_{i=2}^{n}\beta_{1,i}x_i] = \sum_{i\geq 4}(*)x_i.$$

Comparing the coefficients at the corresponding basic elements we get equality $\alpha'_{2,2}\beta'_{2,3} = 0$, i.e. we have a contradiction with supposition $h = 3$.

If $h \geq 4$, then we obtain $\beta'_{h-1,3}\neq 0$. Consider the chain of equalities

$$[y_{h-2}, y_2] = [y_{h-2}, [y_1, x_1]] = [[y_{h-2}, y_1], x_1] - [[y_{h-2}, x_1], y_1] =$$

$$= [\sum_{i=3}^{n}\beta_{h-2,i}x_i, x_1] - [y_{h-1}, y_1] = -\beta_{h-1,3}x_3 + \sum_{i\geq 4}(*)x_i.$$

Since $y_{h-2} \in L^{h-1}$ and $y_2 \in L^3$ then $x_3 \in L^{h+2} = \{x_4, \dots, x_n, y_{h-1}, \dots, y_m\}$, which is a contradiction with the assumption that the nilindex of L is equal to $n + m$. □

Remark 3.1. *In this subsection we used product* $[y_1, x_1] = y_2$*. However, it is not difficult to check that the obtained results are also true under the condition* $[y_1, x_1] = 0$.

3.2. The case of both generators lie in L_1.

Theorem 3.5. *Let* $L = L_0 \oplus L_1$ *be a Leibniz superalgebra from* $Leib_{n,m}$ *with characteristic sequence equal to* $(n_1, \dots, n_k | m_1, \dots, m_s)$*, where* $n_1 \leq n-2, \ m_1 \leq m-1$ *and let both generators lie in* L_1*. Then* L *has a nilindex less than* $n+m$*.*

Proof. Since both generators of the superalgebra L lie in L_1, they are linear combinations of the elements $\{y_1, y_{m_1+1}, \dots, y_{m_1+\dots+m_{s-1}+1}\}$. Without loss of generality we may assume that y_1 and y_{m_1+1} are generators.

Let $L^{2t} = \{x_i, x_{i+1}, \dots, x_n, y_j, \dots, y_m\}$ for some natural number t and let $z \in L$ be an arbitrary element such that $z \in L^{2t} \backslash L^{2t+1}$. Then z is obtained by the products of even number of generators. Hence $z \in L_0$ and $L^{2t+1} = \{x_{i+1}, \dots, x_n, y_j, \dots, y_m\}$. In a similar way, having $L^{2t+1} = \{x_{i+1}, \dots, x_n, y_j, \dots, y_m\}$ we obtain $L^{2t+2} = \{x_{i+1}, \dots, x_n, y_{j+1}, \dots, y_m\}$.

From the above arguments we conclude that $n = m-1$ or $n = m-2$ and

$$L^3 = \{x_2, \dots, x_n, y_2, y_3, \dots, y_{m_1}, y_{m_1+2}, \dots, y_m\}.$$

Applying the above arguments we get that an element of form $B_1 y_2 + B_2 y_{m_1+2} + B_3 y_{m_1+m_2+1}$ disappears in L^4. Moreover, there exist two elements $B_1' y_2 + B_2' y_{m_1+2} + B_3' y_{m_1+m_2+1}$ and $B_1'' y_2 + B_2'' y_{m_1+2} + B_3'' y_{m_1+m_2+1}$ which belong to L^4, where

$$rank \begin{pmatrix} B_1 & B_2 & B_3 \\ B_1' & B_2' & B_3' \\ B_1'' & B_2'' & B_3'' \end{pmatrix} = 3.$$

Since x_2 does not belong to L^5 then the elements $B_1' y_2 + B_2' y_{m_1+2} + B_3' y_{m_1+m_2+1}$, $B_1'' y_2 + B_2'' y_{m_1+2} + B_3'' y_{m_1+m_2+1}$ lie in L^5. Hence, from the notations

$$[x_1, y_1] = \alpha_{1,2}(B_1 y_2 + B_2 y_{m_1+2} + B_3 y_{m_1+m_2+1}) + \alpha_{1,m_1+2}(B_1' y_2 + B_2' y_{m_1+2} + B_3' y_{m_1+m_2+1}) +$$

$$+\alpha_{1,m_1+m_2+1}(B_1'' y_2 + B_2'' y_{m_1+2} + B_3'' y_{m_1+m_2+1}) + \sum_{j=3, j\neq m_1+2, m_1+m_2+1}^{m} \alpha_{1,j} y_j.$$

$$[x_1, y_{m_1+1}] = \delta_{1,2}(B_1 y_2 + B_2 y_{m_1+2} + B_3 y_{m_1+m_2+1}) + \delta_{1,m_1+2}(B_1' y_2 + B_2' y_{m_1+2} + B_3' y_{m_1+m_2+1}) +$$

$$+\delta_{1,m_1+m_2+1}(B_1'' y_2 + B_2'' y_{m_1+2} + B_3'' y_{m_1+m_2+1}) + \sum_{j=3, j\neq m_1+2, m_1+m_2+1}^{m} \delta_{1,j} y_j,$$

we have $(\alpha_{1,2}, \delta_{1,2}) \neq (0,0)$.

Similarly, from the notations

$$[B_1 y_2 + B_2 y_{m_1+2} + B_3 y_{m_1+m_2+1}, y_1] = \beta_{2,2} x_2 + \beta_{2,3} x_3 + \dots + \beta_{2,n} x_n,$$

$$[B_1 y_2 + B_2 y_{m_1+2} + B_3 y_{m_1+m_2+1}, y_{m_1+1}] = \gamma_{2,2} x_2 + \gamma_{2,3} x_3 + \dots + \gamma_{2,n} x_n,$$

we obtain the condition $(\beta_{2,2}, \gamma_{2,2}) \neq (0,0)$.

Consider the product

$$[x_1, [y_1, y_1]] = 2[[x_1, y_1], y_1] = 2\alpha_{1,2}[B_1 y_2 + B_2 y_{m_1+2} + B_3 y_{m_1+m_2+1}, y_1] +$$

$$+2\alpha_{1,m_1+2}[B_1' y_2 + B_2' y_{m_1+2} + B_3' y_{m_1+m_2+1}, y_1] +$$

$$2\alpha_{1,m_1+m_2+1}[B_1'' y_2 + B_2'' y_{m_1+2} + B_3'' y_{m_1+m_2+1}, y_1] +$$

$$+2 \sum_{j=3, j\neq m_1+2, m_1+m_2+1}^{m} \delta_{1,j}[y_j, y_1] = 2\alpha_{1,2}\beta_{2,2} x_2 + \sum_{i\geq 3} (*) x_i.$$

On the other hand,

$$[x_1,[y_1,y_1]] = [x_1, \beta_{1,1}x_1 + \beta_{1,2}x_2 + \dots + \beta_{1,n}x_n] = \sum_{i\geq 3}(*)x_i.$$

Comparing the coefficients at the basic elements in these equations we obtain $\alpha_{1,2}\beta_{2,2} = 0$.

Analogously, considering the product $[x_1,[y_{m_1+1}, y_{m_1+1}]]$, we obtain $\delta_{1,2}\gamma_{2,2} = 0$.

From this equations and the conditions $(\beta_{2,2},\gamma_{2,2}) \neq (0,0)$, $(\alpha_{1,2},\delta_{1,2}) \neq (0,0)$ we easily obtain that the solutions are $\alpha_{1,2}\gamma_{2,2} \neq 0, \beta_{2,2} = \delta_{1,2} = 0$ or $\beta_{2,2}\delta_{1,2} \neq 0, \alpha_{1,2} = \gamma_{2,2} = 0$.

Consider the following product

$$[[x_1,y_1],y_{m_1+1}] = [x_1,[y_1,y_{m_1+1}]] - [[x_1,y_{m_1+1}],y_1] = -\delta_{1,2}\beta_{2,2}x_2 + \sum_{i\geq 3}(*)x_i.$$

On the other hand,

$$[[x_1,y_1],y_{m_1+1}] = \alpha_{1,2}\gamma_{2,2}x_2 + \sum_{i\geq 3}(*)x_i.$$

Comparing the coefficients of the basic elements in these equations we obtain irregular equation $\alpha_{1,2}\gamma_{2,2} = -\beta_{2,2}\delta_{1,2}$. It is a contradiction with supposing the nilindex of the superalgebra equal the $n+m$. And the theorem is proved. □

Thus, the results of the Theorems 3.1–3.5 show that the Leibniz superalgebras with nilindex $n+m$ $(m \neq 0)$ are the superalgebras mentioned in section 2. Hence, we completed the classification of the Leibniz superalgebras with nilindex $n+m$.

References

[1] Albeverio S., Ayupov Sh.A., Omirov B.A. *On nilpotent and simple Leibniz algebras.* Comm. in Algebra, 33(1), 2005, p. 159–172.

[2] Bordemann M., Gómez J.R., Khakimdjanov Yu., Navarro R.M. *Some deformations of nilpotent Lie superalgebras*, J. Geom. and Phys., 57, 2007, p. 1391–1403

[3] Ayupov Sh. A., Khudoyberdiyev A. Kh., Omirov B. A. *The classification of filiform Leibniz superalgebras of nilindex n+m*, Acta Math. Sinica (english series), vol. 25(1), 2009, p. 171–190.

[4] Camacho L.M., Gómez J.R., Navarro R.M., Omirov B.A. *Classification of some nilpotent class of Leibniz superalgebras*, appear in Acta Math. Sinica (english series), 2009, arXiv:math/0611636.

[5] Camacho L.M., Gómez J.R., Omirov B.A, Khudoyberdiyev A.Kh *On complex Leibniz superalgebras of nilindex n+m*, submitted to J. Geom. and Phys., arXiv:math/0812.2156.

[6] Camacho L.M., Gómez J.R., Omirov B.A, Khudoyberdiyev A.Kh. *On the description of Leibniz superalgebras of nilindex* $n+m$, arXiv:0902.2884v1

[7] Gilg M. *Super-algèbres de Lie nilpotentes:* PhD thesis. University of Haute Alsace, 2000. – 126 p.

[8] Gómez J.R., Khakimdjanov Yu., Navarro R.M. *Some problems concerning to nilpotent Lie superalgebras*, J. Geom. and Phys., 51(4), 2004, p. 473–486.

[9] Kac V.G. *Lie superalgebras*, Advances in Math., vol. 26 (1), 1977, p. 8-96.

[10] Livernet M. *Rational homotopy of Leibniz algebras*, Manuscripta Mat., vol. 96, 1998, p. 295–315.

[11] Loday J.-L. *Une version non commutative des algébres de Lie: les algébres de Leibniz*, Ens. Math., 39, 1993, p. 269–293.

[J.R. Gómez] Dpto. Matemática Aplicada I. Universidad de Sevilla. Avda. Reina Mercedes, s/n. 41012 Sevilla. (Spain)

E-mail address: `jrgomez@us.es`

[A.Kh. Khudoyberdiyev – B.A. Omirov] Institute of Mathematics and Information Technologies of Academy of Uzbekistan, 29, F.Hodjaev srt., 100125, Tashkent (Uzbekistan)

E-mail address: `khabror@mail.ru --- omirovb@mail.ru`